



Aquatic Invasive Species Quick Guide



Golden Sands
Resource Conservation
& Development Council, Inc.

Rusty Crayfish (*Orconectes rusticus* Girard)

Description: The rusty crayfish is a large crayfish of the family Cambaridae. Adults can reach six inches in length, including the claws. Most of the body is tan to light brown, but each side of the carapace has a rusty brown spot. Claws are large and typically have brightly colored tips above dark bands. Rusty crayfish are omnivores, feeding primarily on aquatic plants, snails, and other small invertebrates. They can commonly be found hiding under rocks, logs, and other debris. Rusty crayfish typically live 3-4 years.



Rusty crayfish are mostly light brown, with bright claw tips.

North American Distribution: Rusty crayfish are most abundant in the western Great Lakes states, but have been documented from Minnesota and Iowa eastward to Maine, and in northern New Mexico.



Each side of the carapace has a rusty brown spot.

Dispersal Vectors: Rusty crayfish are native to the Ohio River Basin, and were likely transported to the Midwest United States as bait by fishermen. Rusty crayfish quickly colonize lakes and streams by producing several hundred eggs per female each season. Eggs are protected under the female's tail until they hatch.

Ecological Impacts: Rusty crayfish are larger and more aggressive than most native crayfish, and are able to outcompete native species for food and habitat. Rusty crayfish consume large amounts of aquatic invertebrates, small fish, fish eggs, tadpoles, native crayfish, plants, and other aquatic life. They can hasten spread of Eurasian watermilfoil and other aggressive plants by cutting the stems, which then take root elsewhere.

Control Options: Manual trapping is effective for rusty crayfish. Always follow local trapping regulations. Modification of size limits of predator fish species can be effective. Walleye, smallmouth bass, largemouth bass, and yellow perch will consume young rusties. This strategy combined with trapping of large adults can reduce rusty crayfish populations.

An effective, safe pesticide for rusty crayfish has not been found. Although crayfish-selective pesticides exist, they are also harmful to native crayfish species.

An effective biological control agent has not been found. A parasite called *Microphallus* infects rusty crayfish (and other crayfish species), and is currently being researched. Lakes with abundant rusty crayfish are often found to have no *Microphallus* present.



Eggs and newly hatched young are held under the female's tail for protection.

Additional Information:

Hein, C.L., Roth, B.M., Ives, A.R., and M. Jake Vander Zanden. 2006. Fish predation and trapping for rusty crayfish (*Orconectes rusticus*) control: a whole-lake experiment. *Can. J. Fish. Aquat. Sci.* 63: 383-393.

Wisconsin Department of Natural Resources. Potential impacts to rusty crayfish (*Orconectes rusticus*) populations from a parasite, *Microphallus* sp.

Photo Credit: Paul Skawinski